HARVARD UNIVERSITY



LIBRARY

OF THE

Museum of Comparative Zoology





SECOND EDITION.

REMAINS

OF

DOMESTIC ANIMALS

DISCOVERED AMONG

POST-PLEIOCENE FOSSILS

 \mathbb{N}

SOUTH-CAROLINA.

BA

FRANCIS S. HOLMES, A.M.,

Professor of Geology and Zoology Codego of Charleston, S. C., Menner of the American Association for the Advancement of Sciences Corresponding Member Acad. Nat. Sec., Pudadesphas, and Lycenia Nat. History, New York, and Curator of the Museum Nat. History, Charleston, S. C.

ALSO, EXTRACTS FROM A PAPER BY

PROFESSOR LEIDY, OF PHILADELPHIA.

AND A LETTER BY

PROFESSOR AGASSIZ.

CHARLESTON, S. C.

JAMES AND WILLIAMS, PRINTERS,
, Steam Fresses, 16 State Street.

1858.



The Forse not Originally Imported into America from the East.

To the Editors of the Charleston Mercury:

Messes. Editors:—Under the above head you published, a short time since, a notice of the fossil remains discovered in the marl-beds of Ashley River, and included a short extract from a paper written by the distinguished anatomist, Professor Leidy of Phil-

adelphia.

Your notice seems to have attracted general attention, but I regret to find by the letter you kindly sent me, that one of your correspondents is afraid the article in question may cause some misapprehension of the views of Professor Liedy in regard to the origin of the recent or existing species of the horse in North America. I have read with care the letter of your Philadelphia correspondent, and for reasons that will hereafter appear, I must, beg permission to reply to it.

Why should he infer from your article on the fossil remains of the horse, or from the extract from Professor Leidy's paper, that allusion was made to the recent horse? The opinion of Professor Leidy in regard to the origin of our domestic horse, is not alluded to; your remarks are confined to the fossil remains of this animal, and not a word is said about

recent species, or those now in existence,

The caption of the article must have excited the apprehension expressed by the writer, and as this heading may have been suggested by my note to you, inclosing the extract, I feel myself responsible for it.

To avoid all misapprehension of the views of a highly esteemed friend, and of my own also, I have concluded to send you the accompanying additional extracts from the same paper; it was kindly prepared for me by Prof. Leidy, as a contribution to my work on the fossils of South-Carolina. These extracts I give now more cheerfully, as I find some time must elapse before the number which will contain the whole article can be published, with its numerous illustrations.

I have also added a few remarks of my own on

these interesting remains.

Respectfully,

FRANCIS S. HOLMES.

Fossils of the Post-Pleiocene.

The post-pleiocene period is marked in the geological sequence, as that interesting epoch when life upon our globe was manifested in those organic forms, chiefly of the same species, that belong to the historical, or present period, and were obviously designed "from the beginning" to be the cotemporaries and companions of man, who appeared immediately afterwards: "The crowning point of creation."

It is the last formation of the cainozoic or tertiary, the epoch just antecedent to the advent of man upon this earth; a period in which, it may be said, the earth had been finally prepared and made ready for him who was to be formed in the likeness of the Creator, and was to have dominion given him over the "fish of the sea, the fowls of the air, the beasts of the earth, the herb and every creeping thing, yea—over all the earth;" a period that will ever be distinguished as the grand connecting link between the past and the present.

Sir Chas. Lyell, who is considered the best authority on tertiary geology, placed this connection in a striking point of view, having ascertained that ninety-five per cent. of the fossils of the post-pleio-

cene period, are identical with living species.

The crust of the earth, as far as the researches of geologists extend, appears to be separated into strongly marked divisions, that seem to have been formed during four distinct and prolonged periods. These periods, for the sake of convenience, have been named in accordance with the class of animals and vegetables in existence during the formation of each, evidences of which we find preserved in their fossil remains.

The first division has been called Azoic, which signifies without life; the rocks of this age consist of granite, gneiss, etc.; they contain no traces of organic forms, and being the lowest or first formed in the series, and originally in an incandescent state, passed, no doubt, into the condition of rocks, before the creation of animals and plants.

In the second division, the *Palæozoic*, or ancient life, we have indications of the first animals and plants created, not one species of which has outlived the convulsions that separated this period from its successor; and that served, apparently, in each case to mark the close of one period and the commence-

ment of another.

The next in order and third of the series, is named the *Mesozoic* or *middle life*, and its fossils belong to that middle or intermediate class of animals, between the most ancient and recent forms. These also in turn became, like their predecessors extinct, and were succeeded by a later creation.

The next and last age is named the Cainozoic—recent life—and by some authors called the Tertiary, or third grand division, and in which we discover, for the first time, the created forms that are to be the

cotemporaries of man.

These divisions, or grand divisions, as they may be called, are again sub-divided into minor periods or formations; the first, consequently the lowest and oldest in the series of the Cainozoic or Tertiary, is called Eocene, for in it we find representatives of the dawn or commencement of that creation; a few species, two or three per cent. only, having been perpetuated down to the present time.

In the Meiocene, the next in age, a larger number of species are found whose existence is thus prolonged.

In the Peliocene, more recent, a majority of the fossils are of recent species; and at last, in the Post-Pleiocene—the most recent—ninety-five per cent., or nearly all the species, continue to recent times.

Now the evidence herein to be adduced will shew that among the fossils collected in South Carolina from beds of this age—Post-Pleiocene—some of which are exposed at Ashley Ferry, Goose Creek, Stono, John's Island, and other localities, a number have been found apparently belonging to animals having specific characters in common, with recent or living species not considered indigenous to this coun-

try, such as the horse, hog, sheep, ox, etc.

A large collection of fossils from this interesting formation were submitted by me about three years ago, to Professor Leidy, of Philadelphia, the eminent paleontologist, for determination; of these a number were returned with the remark, that they appeared to belong to recent species which had become accidental occupants of the same bed with the true fossils. I held the opposite opinion, and believed that these relics were indeed true fossil remains, as they were obtained not only from the banks and deltas of rivers, but a large number from excavations several feet below the surface, and at a distance from any stream, creek, pond, bog or ravine; and in some cases from excavations below the high sandy land of cotton fields.

In a letter to Dr. Nott and Mr. Gliddon,* dated Feb. 10, 1857, Prof. L. writes:

"Sometime since, Professor F. S. Holmes, of Charleston, submitted for my examination, a collection of fossil bones from a post-pleiocene deposit on Ashley River, S. C. Among remains of the extinct horse, the peccary, mylodon, megatherium, mastodon, hipparion, the tapir, the capabara, the beaver, the musk-rat, etc., were some which I considered as belonging to the dog, the domestic ox, the sheep and the hog. Prof. Holmes observes that these remains were taken from an extensive deposit, in which similar ones exist abundantly, and he further adds, that he cannot conceive that the latter should have become mingled with the former, since the introduction of domestic animals into America by Europeans. It is not improbable that the American continent once had, as part of its fauna, representatives of our domestic animals, which subsequently became extinct—though I am inclined to doubt it; but what we have learned of the extinct American horse, will lead me carefully to investigate the subject."

The opportunity for prosecuting this investigation, to some extent, I had the pleasure of affording Professor Leidy, in March last, a month after the date of the above letter. Dr. Hallowell and himself visited me in Charleston, and I accompanied them to Ashley ferry and Goose creek. The annexed extracts are from a paper of Professor Leidy's on this topic, written after his return home to Philadelphia, and he has also kindly sent me a number of very valuable drawings of fossil horse teeth, and other remains obtained from the Carolina beds.†

"The interesting collection of remains of vertebrated animals, which form the subject of the following pages, for the most part have been submitted to the inspection of the author, by Prof. Holmes and Capt. A. H. Bowman, U. S. A., who collected them from the cocene, post-pleiocene, and recent geological formations,

in the vicinity of Charleston, South-Carolina.

"The collections of these gentlemen consist of a most remarkable intermixture of remains of fishes, reptiles and mammals, of the three periods mentioned; and in many cases perhaps we may err in referring a particular species to a certain formation, more especially in the case of the fishes. The remains usually consist of teeth often well preserved, but frequently in small fragments, more or less water-worn, and most of the fossils are stained brown or block.

* Indigenous races of the earth; p. xix.

[†] Lithographs of these figures will appear in the volume, with Prof. L.'s paper.

"By far the greater portion of the fossil remains are obtained from the post-pleiocene deposit of the Ashley River, about ten miles from Charleston. The country in this locality is composed of a base of whitish eocene marl, containing remains of squalodon—sharks and rays—above which is a stratum of post-pleiocene marl, about one foot in thickness, overlaid by about three feet of sand and earth mould

"The post-pleiocene marl contains great quantities of irregular, water-worn fragments of the eocene marl rock from beneath, mingled with sand, blackened pebbles, water rolled fragments of bones, and more perfect remains of fishes, reptiles and manmals,

belonging to the post-pleiocene and eocene fossils.

"On the shores of the Ashley River, where the post-pleocene and eocene formations are exposed, the fossils are washed from their beds, and become mingled with the remains of recent indigenous and domestic animals, and objects of human art, so that when a collection is made in this locality, it is sometimes difficult to determine whether the animal remains belong to the formations mentioned or not. Generally, however, we have been able to ascertain where the fossils belong, which we have had the opportunity of examining, from the fact that the greater number were obtained from the deposits referred to in digging into them some distance from the Ashley River.

"The collections contain remains of the horse, ox, sheep, hog and dog, which I feel strongly persuaded, with the exception of many of those of the first mentioned animal, are of recent date, and have become mingled with the true fossils of the post-pleiocene and eocene formations, where there have been exposed on the banks of the Ashley River and its tributaries. In regard to the remains of the horse, from the facts stated in the account given of them in the succeeding pages, I think it will be conceded that this animal inhabited the United States during the post-pleiocene period, contemporarily with the mastodon megalonyx, and the great broad fronted bison.

"Many of the mammalian remains are of recent animals, or at least are undistinguishable from the corresponding parts of the latter; and if they are not accidental occupants of the post-plei-ocene deposit, are highly interesting, as indicating their contemporaneous existence with many species and genera now ex-

tinet.*

"It appears to be quite well authenticated that the horse, which is now so extensively distributed, both in a wild and domestic condition, throughout North and South America, did not inhabit these continents at the time of their discovery by Europeans. With this fact in view, in conjunction with the circumstance that animal remains of late periods may become accidental

^{*} Remains of the Tapir, Peccary and Cabybara present a similar association of life to that now confined to South America.

occupants of earlier geological formations, we should require strong evidence to be advanced before it is admitted that the Horse belonged to an ancient fauna of the western world. the present time the evidence appears to be sufficiently ample to justify the latter conclusion, and it is further sustained by the discovery, in the same part of the world, of the remains of two

species of the closely allied genus Hipparion.

"Remains of the Horse, discovered in Brazil, Buenos-Ayres Chili, have been indicated by Dr. Lund, Prof. Owen, M. Weddell, and M. Gervais. These remains exhibit no well marked characters distinguishing them from corresponding portions of the skeleton of the recent Horse, and from a comparison of the figures and descriptions which have been given of most of them, together with some remarks of the latter author, it is doubtful whether they belong to more than a single species, the Equus neogens of Dr. Lund.

" Prof. Buckland and Sir John Richardson have described remains of the Horse, discovered in association with those of the Elephant, Moose, Reindeer, and Musk Ox, in the ice cliffs of

Eschscholtz Bay, Arctic America.

"In the United States, remains of the Horse, chiefly consisting of teeth, have been noticed by Drs. Mitchell,* Harlan,† and De Kay, t but these gentlemen have neither given descriptions nor Some of the latter figures by which to identify the specimens. are stated to have been found in the vicinity of Neversink Hills, New Jersey; others in the excavation for the Chesapeake and Ohio Canal, near Georgetown, District of Columbia; and some in the later tertiary deposite on the Neuse River, in the vicinity of Newbern, North Carolina. Dr. DeKay, in speaking of such remains, says, "they resemble those of the common Horse, but from their size apparently belonged to a larger animal," and he refers them to a species with the name of Equus major.

"Dr. R. W. Gibbes has given information of the discovery of teeth of the Horse in the pleiocene deposit of Darlington, South-Carolina; in Richland District of the same State; in Skidaway Island, Georgia, and on the banks of the Potomac river. He further observes that he obtained the tooth of a Horse, from eocene marl, in the Ashley river, South Carolina, but the researches of Prof. Holmes | indubitably indicate the specimen to

have been an accidental occupant of the formation.

"Specimens of isolated teeth, and a few bones of the horse, from the post-pleiocene and recent deposits of this country, have frequently been submitted to my inspection. Many of these I have unhesitatingly pronounced to be relics of the domestic

|| Ibidem, 68.

^{*} Catalogue of Organic Remains, 1826, 7, 8. † Med. a. Phys. Researches, 1835, 267. ‡ Zoology, New York, pl. 1, Mammalia, 108. § Proc. Amer. Assoc; 1850, 66.

horse, though I feel persuaded that many remains of an extinct

species are undistinguishable from the recent one.

"Whether more than one extinct species is indicated among the numerous specimens of teeth I have had the opportunity of examining, I have been unable satisfactorily to determine. The specimens present so much difference in condition of preservation, or change in structure; so much variation in size, from that of the more ordinary horse to the largest English dray horse; and such variableness in constitution, from that of the recent horse to the most complex condition belonging to any extinct species described, that it would be about as easy to indicate a half dozen species as it would two.

Under the circumstances, I would characterize the extinct horse of the United States, as having had about the same size as the recent one, ranging from the more ordinary varieties to the English dray horse, with molar teeth, frequently comparatively simple in construction, but with a strong disposition to

become complex.

"Among the number of teeth of the horse in Prof. Holmes' collection, labelled as coming from the post-pleiocene deposit of Ashley River, there are several, which, from their size, construction and condition of preservation, I feel convinced are of recent date; and these no doubt became mingled with the true fossils of that formation where it is exposed on the Ashley River, in which position I personally found undoubted remains of the recent horse and other domestic animals, and objects of human art, mingled with remains of fishes, reptiles, and mammals, washed by the river from the banks, composed of eocene and post-pleiocene deposits.

"Teeth of an extinct species of Horse, however, undoubtedly belong as true fossils to the post-pleiocene formations in the vicinity of Charleston. These are usually hard in texture, stained brown or black from the infiltration of oxide of iron, sometimes well preserved, but more frequently in a fragmentary condition and water worn. Generally they are not larger than the teeth of the more ordinary varieties of the domestic horse, and sometimes are quite as simple in the plication of their enamel, but usually are more complex and sometimes exceed-

ingly so.

"Figure 1 represents a first superior molar tooth, neither larger nor more complex in structure than the corresponding tooth of the recent Horse. This specimen, which is dense and jet black in color, was obtained by Prof. Holmes from a stratum of ferruginous sand, two inches thick, exposed on the side of a bluff, on Goose Creek, about twelve miles from Charleston.

"Having expressed a desire to see the locality from which the tooth just mentioned was obtained, Prof. Holmes afforded me the opportunity of doing so. The bluff is about thirty feet high; its base is formed of a pleiocene limestone, about fifteen feet thick and composed of the debris of marine shells; above this is the stratum of ferruginous sand, of post-pleiocene age, containing numerous pebbles and rolled fragments of bone all blackened like the tooth obtained from the same position. Overylying the latter stratum, there is a layer of stiff blue clay, about two feet in thickness, and above this there are about twelve feet of sand and earth-mould.

"A similar blackened tooth was obtained from the same for-

mation at Doctor's Swamp, John's Island.

"Figure 4 represents a very remarkably well preserved specimen of a lower molar above referred to, from Georgia, where it was discovered by J. H. Couper, in association with equally well preserved remains of other extinct animals. The tooth is brown in color; and it neither differs in size nor form from its homo-

logue in the recent Horse.

"In the collection of fossils of Prof. Holmes, there is the specimen of an upper first large molar, labelled from Texas, represented in figure 5. The tooth is of the largest comparative size, and exhibits the highest degree of complexity in the folding of its enamel; in both of which characters it differs in such a remarkable degree from the corresponding tooth, represented in figure 5, from the post-pleiocene formation of South-Carolina, that it appears hardly possible that these two teeth should belong to the same species of horse.

"A remarkably well preserved specimen of an upper molar tooth, jet black in color, and an incisor, yellow and quite friable in texture, both belonging to the extinct horse, from North-Carolina, have been submitted to my inspection by Prof. Emmons.

"Among the most interesting of the fossils discovered by Prof. Holmes, in the post-pleiocene beds of the Ashley River, are two molar teeth of a species of the equine genus *Hippotherium*. These are the first remains of the latter discovered in America, and they indicate the smallest known species.

"Both specimens are from the upper jaw; and they are well characterized, not only by the isolation of the internal median enamal column, but also by the complex plication of the interior

or central enamel columns.

"The larger specimen is firm in texture; has the enamel stain-

ed jet-black, and the dentine and cement gray.

"I have personally had the opportunity of inspecting remains of the tapir, found in Texas, Louisiana, Kentucky, Mississippi, Indiana, Ohio and South-Carolina, proving an extensive range of this animal at one time over the country of the United States.

"The specimens which were presented by Dr. Carpenter to the Academy of Natural Sciences of Philadelphia, on close comparison are not found to differ from the corresponding parts of the living tapirus americanus.

"The post-pleiocene deposit of the Ashley River, contain a number of small fragments of molar teeth, and one nearly entire and unworn crown of a second lower molar, which have the same characters of form and size, as in the living tapir. Besides these, the same collections contain fragments of lower molars, and two nearly entire crowns of upper molars, having the exact form of the corresponding teeth of the *T. americanus*, but larger in size.

"Teeth of the beaver, jet black in color, have likewise been obtained from the post-pleiocene deposit of Ashley River.

"The collections contain numerous specimens of blackened molar teeth, together with a few incisors and fragments of jaws, from the Ashley post-pleiocene deposit, which neither differ in form nor size from the corresponding parts of the recent muskrat.

"Remains of Lepus sylvaticus—common gray rabbit—have been found in association with those of other rodents and of the extinct peccary near Galena, Illinois. A few specimens of molar teeth, black in color, apparently belonging to this species, were obtained from the post-pleiocene beds of the Ashley River.

"Several small fragments of teeth of the Megatherium, in Prof. Holmes' collection, were obtained from the post-pleiocene bed of the Ashley River. Previously to the discovery of these specimens, remains of the Megatherium had been found in no other locality of North America, than in the State of Georgia.

"Two small fragments of lower molar teeth of Mylodon Harlani, were obtained from the Ashley post-pleiocene beds. One of the fragments is represented in figures 21, plate xvi, of 'A memoir on the extinct Sloth Tribe of North America,' by the author."

As regards the specimens of human art found as above, it must be remarked that it is only at this locality—Ashley Ferry—that we find such relics. Here at the base of a low bluff, is a beach of Eocene marl; above the bluff is a farm-yard, and all the sweepings of the premises, consisting in part of old hoes, broken plough-shares, and fragments of crockery-ware, etc., are thrown into the river, and lie mingled with the fossils which are washed out of the bluff, and scattered over the surface of the beach telow, which is exposed at low tide. At no other locality on this river, and there are several, viz: Ramsay's, Clement's, Greer's, Middleton's, etc., where similar fossils are found, do we obtain relics of human art; at least, I have never found such.

The fossils from Ashley Ferry present, as a group, the same appearance as those procured inland at some distance from the river, by digging from three to five feet below the surface. Many specimens from the ferry were considered as recent by Professor Leidy; they appear quite fresh and unchanged in color, and their texture not in the slightest degree altered. To one familiar with the fossils of the South Carolina Post-Pleiocene, this excites no surprise, as it is of common occurrence, more especially among the shells; for example, the olive shell—Oliva literata—is found as fresh and highly polished as the recent ones from the sea-beaches along the coast; and Cardium magnum retains often, the delicate yellow and brown markings; common to the species.

The color or texture of a fossil, therefore, does not always absolutely determine its relative age; as Professor Leidy has himself remarked in a foot-note to

his letter alluded to above, viz:

"Fossilization, petrification, or lapidification, is no positive

indication of the relative age of organic remains.

"The Cabinet of the Academy of Natural Sciences, of Philadelphia, contains bones of the megalonyx, and of the extinct peccary, that are entirely unchanged; not a particle of gelatin has been lost, nor a particle of mineral matter added, and, indeed, some of the bones of the former even have portions of articular cartilage and tendinous attachments, well preserved."*

From the foregoing it would apppear that of the ancient fauna of America, which included representatives of many of our present domestic animals, some species have undoubtedly become extinct; but I confess I am not yet prepared to admit from any evidence yet adduced, or from my own examinations that, all of the living species are distinct from those found fossil in the post-pleiocene. The teeth and bones of the rabbit, raccoon, opossum, deer, elk, hog, dog, sheep, ox and horse are often found in these beds, and though associated with those known to be

^{*} Indigenous races of the earth; p. xix.

extinct, such as mastodon, megatherium, hipparion, etc., need not necessarily be referred to extinct races also; since their remains cannot be distinguished

from the bones and teeth of the living species.

Of the mollusca from the same beds about ninety-five per cent are to my mind identically the same with species now living on the coast of South-Carolina. Two species of these shells though extinct, or not in existence here, are now living in numbers on the coast of Florida and the northern shores of the gulf of Mexico;* and two have no living representatives that we can discover.†

The question therefore naturally suggests itself—are the living horses, dogs, hogs, raccoons, opossums, deer, elk, tapirs, beavers, etc., and the one hundred and fifty living shells of the coast, the descendants of the animals whose remains we find fossil in the above

named beds.

It has been just remarked that about ninety-five per cent., or nearly all of the one hundred and fifty shells of molluscous animals from these beds are specifically identical with the recent or living species of the coast,—two are found only at the south of this, and two are extinct. Of the vertebrates from the same bed, the tapir, peccary, raccoon, opossum, deer, musk-rat, rabbit, beaver, and elk have still their living representatives, generically, if not specifically; and even of the identity of species there seems to be no doubt, as no anatomical differences can be discerned. Two of these species, like the mollusca just alluded to, no longer live in South Carolina; the tapir and peccary are only found in South America and Mexico; the musk-rat, elk and beaver, though extinct on the Atlantic coast, are still living in the interior of the country. And though it has been acknowledged that the mastodon, megatherium, elephant, glyptodon, and two species of Equine genera,

† Hyalea. Tellina.

^{*} Strombus pugilis; Gnathodon cuneatum.

etc., are entirely extinct, yet the discoveries made of the remains even of some of these, would indicate that they still existed at a period so recent, that, in the language of Professor Leidy, "it is probable the

red man witnessed their declining existence."

The peccary, or Mexican hog, an animal common in Mexico, is not indigenous to the Atlantic United States; but his bones have been found associated with human remains in caves used as cemeteries by the Aboriginees.* "A tomb in the city of Mexico," according to Clavigero, (?) + "was found to contain the bones of an entire mammoth, the sepulchre appearing to have been formed expressly for their reception." And "Mr. Latrobe relates that during the prosecution of some excavations, near the city of Tezcuco, one of the ancient roads or causeways was discovered, and on one side, only three feet below the surface, in what may have been the ditch of the road, there lay the entire skeleton of a mastodon. It bore every appearance of having been coeval with the period when the road was used."

Again I extract from Prof. Leidy's letter:‡

"The early existence of the genera to which our domestic animals belong, has been adduced as presumptive evidence of the advent of man at a more remote period than is usually assigned. It must be remembered, however, even at the present time that of some of these genera only a few species are domesticated: thus of the existing six species of Equus (Horse) only two have ever been freely brought under the dominion of man.

"The Horse did not exist in America at the time of its discovery by Europeans; but its remains, consisting chiefly of molar teeth, have now been so frequently found in association with those of extinct animals, that it is generally admitted once to have been an aboriginal inhabitant. When I first saw examples of these remains I was not disposed to view them as relics of an extinct species; for although some presented characteristic differences from those of previously known species, others were undistinguishable from the corresponding parts of the domestic horse, and among them were intermediate varieties of form and

^{*} Bradford's American Antiquities, p. 31. † Bradford's American Antiquities, p. 227.

Note and Gliddon, Indigenous races of the earth; p. xviii.

size. The subsequent discovery of the remains of two species of the closely allied extinct genus Hipparion, in addition to the discovery of remains of two extinct equine genera of an earlier geological period, leaves no room to doubt the former existence of the Horse on the American continent, contemporaneously with the Mastodon and Megalonyx: and man probably was his companion."

The result of the whole seems to be, that of the animals found fossil in the post-pleiocene beds, all the mollusca of the present day are undoubtedly a perpetuation of the same species; that of the higher order of vertebrata, the tapir, peccary, raccoon, opossum, deer, elk, and musk rat are equally entitled to be considered the descendants of this ancient race. And if the claims of the mollusca to this distinction rests upon a secure basis, because they are peculiar to this country, and not obnoxious to suspicion of foreign immigration, it must be recollected that this is equally true of the above named animals.

Those which have hitherto been regarded as of recent and European origin, are the horse, sheep, hog and ox; and it must be reserved perhaps for future consideration to determine how far the negative proof of the non-existence of these animals in the country at the time of its discovery may be regarded in each individual case sufficiently strong to settle the question of his extinction and re-introduction, when so many of his associates and contemporaries have succeeded in maintaining an unbroken line of descent down to the present day.

By the steamer Isabel, just arrived from Key West, (March 1st) I received the annexed letter from the learned Professor Agassiz, to whom I had the pleasure of exhibiting these interesting specimens, only a few days ago, when he passed through Charleston, en route for Key West:

KEY WEST, FEB. 25th, 1858.

Professor F. S. Holmes:

My Dear Sir:—I have not forgotten my promise to write you my impressions respecting your important discoveries of fossil mammalia in the post-pieiocene beds of South-Carolina. Indeed I have been thinking of them continually since I saw them, and nothing impressed me so deeply for many years past as the sight of these bones. I consider their careful study in all their relations as of the utmost importance for the progress of .our science. It is true there is hardly any thing of interest in the animals themselves, since they appear to be all well known types, but their simultaneous occurrence in the same beds, showing that they have lived together at a time when the white man had not yet planted himself upon this continent, render their association as undisputed. How does it happen, that horses, sheep, bulls and hogs, not distinguishable from our domestic species existed upon this continent, together with the deer, the musk-rat, the beaver, the hare, the opossom, the tapir, which in our days are peculiar to this continent, and not found in the countries where our domesticated animals originated? whole matter might seem to admit of an easy solution by supposing that the native American horse, sheep, bull, and hog were different species from those of the old world, even though the parts preserved show no specific differences; but this would be a mere theoretical solution of a difficulty which seems to me to have far deeper meaning, and to bear directly upon the question of the first origin of organized beings.

The circumstances under which these remains are found, admit of no doubt but the animals from which they are derived, existed in North America long before this continent was settled by the white race of men, together with animals which to this day are common in the same localities, such as the deer, the musk-rat, the opossum and others only now found in South America, such as the tapir. This shows beyond the possibility of a controversy, that animals which cannot be distinguished from one another, may originate independently in different fauna, and I take it that the facts you have brought together, are a satisfactory proof that horses, sheep, bulls and hogs not distinguishable at present from the domesticated species, were called into exisience upon the continent of North America prior to the coming of the white race to these parts, and that they had already disappeared here when the new comers set foot upon this continent; but the presence of tapir teeth among the rest show also that a genus peculiar to South America and the Sunda Islands existed also in North America in those days, and that its representative of that period is not distinguishable from the South

American species.

It would be desirable in this stage of the enquiry to compare your tapir teeth with those of the species from Central America, which is considered distinct from the Brazilian species. This circumstance leads naturally to the question of the specific identity of all these animals with those now living in the same locality, and with the domesticated species. And here I confess the difficulty to be almost insuperable, or at least hardly approachable in the present state of our science, when the views of naturalists are so divided as to what are species among the genera bos, ovis, capra. For myself, I entertain doubt respecting the unity of origin of the domesticated horses. But whatever be the final result of this enquiry, this much is already established by the fossils you have collected, that horses, hogs, bulls and sheep were among the native animals of North America, as early as the common American deer, the opossum, the beaver, the musk rat, etc. What remains to be settled respecting their specific identity is involved in the controversy now carried on between naturalists, who admit specific distinctions upon a very wide range of differences, and those who limit them within narrow boundaries. But the final solution of this point can in no way lessen the interest of your discoveries.

Should you publish anything upon this subject, let me have your notice, for I am deeply interested in the subject, as I always

shall be, in everything you do.

Ever truly your friend, L. AGASSIZ.


CATALOGUE OF MAMMALIA FROM THE POST-PLEIO-CENE.

EXTINCT Species.—Mastodon, Megatherium, Megalonyx, Glyptodon, Mylodon and Hipparion, 2 species.


Not now Found on the Atlantic Coast, but Indigenous to North America.—Bison, Tapir, Peccary, Beaver, Musk rat, and Elk.

THE DEER, RACCOON, OPOSSUM, RABBIT AND THE FOLLOWING DOMESTIC ANIMALS—Horse, Hog, Sheep, Dog and Ox are not distinguishable from the living species.









3 2044 107 335 556

No. land